



# HUMAN CYTOMEGALOVIRUS AND THE CNS: IMMUNITY AND INFLAMMATION

C. Cipriani<sup>1</sup>, X. Borshi<sup>2</sup>, M. Gulino<sup>3</sup>, M.T. Miele<sup>1\*</sup>

<sup>1</sup>Department of Experimental Medicine, Medical School, University of Rome “Tor Vergata”, Rome, Italy;

<sup>2</sup>School of Specialization in Clinical Pathology and Clinical Biochemistry, Medical School, University of Rome “Tor Vergata”, Rome, Italy;

<sup>3</sup>Department of Clinical Science and Translational Medicine, Medical School, University of Rome “Tor Vergata”, Rome, Italy

\*Correspondence to:

Dr Martino Tony Miele

e-mail: [miele@med.uniroma2.it](mailto:miele@med.uniroma2.it)

## ABSTRACT

**Human cytomegalovirus (HCMV) is a highly widespread  $\beta$ -herpes virus that affects all age groups and is largely asymptomatic. The virus is transmitted through contact with bodily fluids and affects the central nervous system (CNS), causing neuropsychiatric disorders and contributing to immune activation and inflammation. HCMV infection can cause significant morbidity and be life-threatening for immunocompromised individuals and those taking immunosuppressant drugs. The virus can cause viral encephalitis in the immunocompromised, frequently resulting in brain damage. In addition, HCMV can cause serious infections affecting newborns with severe symptoms. In newborns, the virus can cross the blood-brain barrier (BBB), infecting brain cells and cerebrospinal fluid. Furthermore, HCMV can infect microglia, immune cells of the CNS with functions similar to macrophages, by activating toll-like receptors (TLRs) with the release of pro-inflammatory cytokines, including IL-1, TNF, and IL-6. This process leads to inflammation and neuropathic pain, which can be reduced by inhibiting these cytokines. Infection can also affect neurons and contribute to glioma and other brain pathologies. Here, the pathogenesis of HCMV infection involving the immune system and the effects on the CNS will be reported.**

**Keywords:** *cytomegalovirus, CNS, immunity, inflammation, microglia; infection*

## INTRODUCTION

Human cytomegalovirus (HCMV) is a  $\beta$ -herpes virus that is widespread worldwide and primarily asymptomatic since the immune system contains it. HCMV is a virus that can affect any age group, with an incidence of about 50%-60% in developed countries (1, 2), contributing to immune activation and inflammation (3). In addition, HCMV can affect the central nervous system (CNS), causing neuropsychiatric disorders (4) and the immune system, particularly in immunocompromised individuals and those taking immunosuppressant drugs (5, 6).

The virus is transmitted from an infected individual to a healthy individual through contact with body fluids, including breast milk, saliva, urine, blood, sperm, tears, wounds, and pus. (7). HCMV infection can remain permanently latent, and its reactivation in subjects with immune pathologies can cause serious damage to vital organs (8). HCMV

Received: 23 September, 2022

Accepted: 28 October, 2022

2279-5855 (2022)

Copyright © by BIOLIFE 2022

This publication and/or article is for individual use only and may not be further

reproduced without written permission from the copyright holder.

Unauthorized reproduction may result in financial and other penalties

Disclosure: all authors report no conflicts of interest relevant to this article.

infection appears to affect immunodeficient patients but can also affect individuals with an efficient immune system. Therefore, HCMV infections can have different degrees of severity. It is one of the most frequent serious infections affecting newborns, estimated to be the leading cause of congenital infection in developed countries with an incidence between 0,3% and 2,3% of all births, presenting with cranial bone changes, mental retardation, microcephaly, jaundice, hearing loss, skin lesions, impaired neurodevelopment, and other symptoms (9,10). This infection most commonly affects children with Down syndrome (trisomy 21). HCMV is also transmitted in bodily fluids from infected but symptom-free individuals (11). Most newborns can render the infection asymptomatic, and the severity of the disease depends on the individual immune response. HCMV infections can be lethal, especially in immunosuppressed patients with impaired T lymphocyte response. In the most severe pathological cases, the CNS and the hematopoietic and respiratory systems can be altered, especially in the first months of the newborn's life (12).

### ***HCMV Immunity and Inflammation***

HCMV, present in body fluids, is primarily transported by myeloid cells and mainly affects certain types of cells, such as endothelial, epithelial, smooth muscle, and fibroblast cells (13). HCMV is transmitted from mother to child across the placenta, passes to the fetus, enters the bloodstream, and reaches the brain (14); the infection mainly affects newborns, while adults show a higher resistance. Since the newborn blood-brain barrier (BBB) is not yet formed, the virus can infect the brain cells and cerebrospinal fluid (15). The virus initially targets and replicates in astrocytes, cells important in synapse formation and supporting the BBB (12).


HCMV can cause viral encephalitis, a very common disease after brain infection, responsible for permanent CNS damage, which can even lead to patient death (16). Viral encephalitis can be caused by various infections mediated by the immune cells responsible for the inflammation. HCMV encephalitis affects most patients with compromised immune systems, and the disease can be diagnosed by polymerase chain reaction (PCR) analysis. Brain damage caused by HCMV is relatively frequent and, in certain circumstances, can cause IL-1beta-mediated fever, chills, fatigue syndrome, myalgia, motor deficits with paraplegia, hypoesthesia, paraesthesia, dysesthesia, anaesthesia, disorientation, brain fog, bilateral visual loss, urinary retention, constipation, and coma (Table I).

**Table I.** *Infections caused by HCMV*

<b><i>Most frequent manifestations:</i></b>	1) colitis	2) meningitis	3) encephalitis	4) myelitis
<b><i>Pathological states of other organs:</i></b>	1) hemolytic anemia	2) thrombocytopenia	3) venous and/or arterial vascular thrombosis	
	4) uveitis	5) pneumonia	6) splenic rupture	
	7) coagulation	8) pancytopenia	9) myelodysplastic changes	
	10) abdominal pain	11) headache	12) jaundice	

Microglia are long-lived immune cells in the brain that viruses, including HCMV, can activate. The innate immune defence of these cells protects against infection in the CNS, although in the case of HCMV, microglia are not the primary target. Microglia perform a function similar to peripheral blood monocytes in the CNS, protecting the brain from infection and inducing inflammation by releasing inflammatory mediators (17). Therefore, these cells appear to be not easily infected by HCMV. However, the virus can infect microglia by activating toll-like receptors (TLRs) involved in glial activation (18). It binds to TLR4, activating pro-inflammatory cytokines such as IL-1 beta, TNF-alpha and IL-6, causing inflammation and neuropathic pain (19, 20) (Fig. 1). These cytokines generated primarily by microglia, are the most studied inflammatory mediators in CNS insults, as several activators including viruses upregulate them. Their inhibition significantly reduces the inflammatory and neuropathic pain induced by pathogenic viruses. However, the exact mechanism(s) remain to be confirmed (21).

Neurons, which have little possibility of replication, can be weakly infected by viruses which exploit neuronal metabolism to replicate and induce apoptosis (22). In addition, neuronal stem cells are also resistant to HCMV infection. However, the ependymal glial cells of the CNS have cilia that move the cerebrospinal fluid of the CNS and are easily infected by the virus (12).



**Fig. 1.** Here, we depict inflammation induced by HCMV in the brain, which provokes the generation of inflammatory cytokines and neurotransmitters such as substances P and neurotensin, leading to inflammation.

An immunodeficiency involving CD8+ and/or CD4+ T cells is important for the reactivation of HCMV infection, an effect to be confirmed by future studies (23). Infection with HCMV can also promote the onset of tumors such as glioma and other brain pathologies (24). In addition, virus infection in the brain attracts T cells, exhibiting an immune and inflammatory effect. In some cases, specific antivirals can relieve symptoms, including systemic inflammation, and avoid death. (25, 26).

However, the pathogenesis of HCMV infection is still unclear, and therefore, future studies, especially involving the immune system, are needed to unravel the many obscure points that arise in the CNS.

#### *Conflict of interest*

The authors declare that they have no conflict of interest.

#### **REFERENCES**

1. Cytomegalovirus (CMV) and Congenital CMV Infection. [(accessed on 10 October 2022)]; Available online: <https://www.cdc.gov/cmV/overview.html>
2. Zuhair M, Smit GSA, Wallis G, et al. Estimation of the worldwide seroprevalence of cytomegalovirus: A systematic review and meta-analysis. *Reviews in Medical Virology*. 2019;29(3):e2034. doi:10.1002/rmv.2034
3. van de Berg Pablo J, Heutinck Kirstin M, Raabe R, et al. Human Cytomegalovirus Induces Systemic Immune Activation Characterized by a Type 1 Cytokine Signature. *The Journal of Infectious Diseases*. 2010;202(5):690-699. doi:10.1086/655472
4. Zhang XY, Fang F. Congenital human cytomegalovirus infection and neurologic diseases in newborns. *Chinese Medical Journal*. 2019;132(17):2109-2118. doi:10.1097/cm9.0000000000000404
5. Griffiths P, Reeves M. Pathogenesis of human cytomegalovirus in the immunocompromised host. *Nature Reviews Microbiology*. 2021;19(12):759-773. doi:10.1038/s41579-021-00582-z
6. Griffiths PD. Cytomegalovirus infection complicating immunosuppressive treatment. *Postgraduate Medical Journal*. 1988;64(747):1-3. doi:10.1136/pgmj.64.747.1

7. Cannon MJ, Schmid DS, Hyde TB. Review of cytomegalovirus seroprevalence and demographic characteristics associated with infection. *Reviews in Medical Virology*. 2010;20(4):202-213. doi:10.1002/rmv.655
8. Forte E, Zhang Z, Thorp EB, Hummel M. Cytomegalovirus Latency and Reactivation: An Intricate Interplay With the Host Immune Response. *Frontiers in Cellular and Infection Microbiology*. 2020;10. doi:10.3389/fcimb.2020.00130
9. Nassetta L, Kimberlin D, Whitley R. Treatment of congenital cytomegalovirus infection: implications for future therapeutic strategies. *Journal of Antimicrobial Chemotherapy*. 2009;63(5):862-867. doi:10.1093/jac/dkp083
10. Gulino M, Miele MT, Marcuccilli F, Cammarano A, Montanari Vergallo G. Late termination of pregnancy in case of congenital CMV infection: ethics, medicine and law. *J Matern Fetal Neonatal Med*. 2022 Dec;35(25):9786-9791. doi: 10.1080/14767058.2022.2053104.
11. Schottstedt V, Blümel J, Burger R, et al. Human Cytomegalovirus (HCMV) – Revised. *Transfusion Medicine and Hemotherapy*. 2010;37(6):365-375. doi:10.1159/000322141
12. Krstanović F, Britt WJ, Jonjić S, Brizić I. Cytomegalovirus Infection and Inflammation in Developing Brain. *Viruses*. 2021;13(6):1078. doi:10.3390/v13061078
13. Aguiar A, Galinato M, Bradley Silva M, Toth B, McVoy MA, Hertel L. Human Cytomegalovirus Replication and Infection-Induced Syncytia Formation in Labial, Foreskin, and Fetal Lung Fibroblasts. *Viruses*. 2021;13(12):2355. doi:10.3390/v13122355
14. Buxmann H, Hamprecht K, Meyer-Wittkopf M, Friese K. Primary Human Cytomegalovirus (HCMV) Infection in Pregnancy. *Deutsches Ärzteblatt Online*. 2017;114(4):45-52. doi:10.3238/arztebl.2017.0045
15. Alcendor DJ, Charest AM, Zhu WQ, Vigil HE, Knobel SM. Infection and upregulation of pro-inflammatory cytokines in human brain vascular pericytes by human cytomegalovirus. *Journal of Neuroinflammation*. 2012;9(1). doi:10.1186/1742-2094-9-95
16. Slavuljica I, Kvešták D, Csaba Huszthy P, Kosmac K, Britt WJ, Jonjić S. Immunobiology of congenital cytomegalovirus infection of the central nervous system—the murine cytomegalovirus model. *Cellular and Molecular Immunology*. 2015;12(2):180-191. doi:10.1038/cmi.2014.51
17. Bachiller S, Jiménez-Ferrer I, Paulus A, et al. Microglia in Neurological Diseases: A Road Map to Brain-Disease Dependent-Inflammatory Response. *Frontiers in Cellular Neuroscience*. 2018;12(488). doi:10.3389/fncel.2018.00488
18. Patrycy M, Chodkowski M, Krzywowska M. Role of Microglia in Herpesvirus-Related Neuroinflammation and Neurodegeneration. *Pathogens*. 2022;11(7):809. doi:10.3390/pathogens11070809
19. Compton T, Kurt-Jones EA, Boehme KW, et al. Human Cytomegalovirus Activates Inflammatory Cytokine Responses via CD14 and Toll-Like Receptor 2. *Journal of Virology*. 2003;77(8):4588-4596. doi:10.1128/jvi.77.8.4588-4596.2003
20. Jarvis MA, Borton JA, Keech AM, et al. Human Cytomegalovirus Attenuates Interleukin-1 $\beta$  and Tumor Necrosis Factor Alpha Proinflammatory Signaling by Inhibition of NF- $\kappa$ B Activation. *Journal of Virology*. 2006;80(11):5588-5598. doi:10.1128/jvi.00060-06
21. Tang XD, Ji TT, Dong JR, et al. Pathogenesis and Treatment of Cytokine Storm Induced by Infectious Diseases. *International Journal of Molecular Sciences*. 2021;22(23):13009. doi:10.3390/ijms222313009
22. Allsopp TE, Scallan MF, Williams A, Fazakerley JK. Virus infection induces neuronal apoptosis: A comparison with trophic factor withdrawal. *Cell Death & Differentiation*. 1998;5(1):50-59. doi:10.1038/sj.cdd.4400298
23. Brizić I, Šušak B, Arapović M, et al. Brain-resident memory CD8<sup>+</sup>T cells induced by congenital CMV infection prevent brain pathology and virus reactivation. *European Journal of Immunology*. 2018;48(6):950-964. doi:10.1002/eji.201847526
24. Hu M, Wang B, Qian D, et al. Human cytomegalovirus immediate-early protein promotes survival of glioma cells through interacting and acetylating ATF5. *Oncotarget*. 2017;8(19):32157-32170. doi:10.18632/oncotarget.17150
25. Devetag FC, Boscaroli L. Cytomegalovirus Meningoencephalitis with Paroxysmal Course in Immunocompetent Adults: A New Nosographical Entity. *European Neurology*. 2000;44(4):242-247. doi:10.1159/000008244
26. Eddleston M, Peacock S, Juniper M, Warrell DA. Severe Cytomegalovirus Infection in Immunocompetent Patients. *Clinical Infectious Diseases*. 1997;24(1):52-56. doi:10.1093/clinids/24.1.52