



VITAMIN E ACTIVITY IN THE CENTRAL NERVOUS SYSTEM

F. Mastrangelo^{1*}

¹Oral Surgery Specialist, Associate Professor, Oral Surgery and Implantology, School of Dentistry, Clinical and Experimental Medicine Dept., University of Foggia, Foggia, Italy

*Correspondence to:

Filiberto Mastrangelo, MD
Oral Surgery Specialist, Associate Professor,
Oral Surgery and Implantology, School of Dentistry,
Clinical and Experimental Medicine Dept.,
University of Foggia, Foggia, Italy
e- mail: filibertomastrangelo@hotmail.com

KEYWORDS: *Vitamin E, anti-oxidant, brain, alpha-tocopherol, Alzheimer's disease, Parkinson's disease*

Nutrients are necessary for the normal physiological function of humans and animals. Without the supply of Vitamin E and mineral salts, no metabolic process in our body, including the brain, could work (1). Vitamins, identified for the first time in 1912, are organic compounds necessary for good health and lifestyle (2). However, long before it was known, foods were necessary for animal life. To date, 13 vitamins can be synthesised in a laboratory and introduced in a regular diet, although our body produces some vitamins (3). Vitamins are classified into two large groups: water-soluble and fat-soluble (4). The fat-soluble vitamins are vitamin A or retinol; vitamin D or calciferol; vitamin E or tocopherol; and vitamin K (5). Water-soluble vitamins are: Vitamin B1 or thiamine; Vitamin B2 or riboflavin; vitamin B3 or PP and niacin; vitamin B5 or pantothenic acid; vitamin B6 or pyridoxine; vitamin B8 or H or biotin; vitamin B9 or folic acid; vitamin B12 or cobalamin, and vitamin C or ascorbic acid (6, 7). Vitamin E is considered the primary fat-soluble anti-oxidant of all vitamins and is involved in lipid peroxidation damage (8-10). Peroxidation is a reaction involved in ageing processes and other neurodegenerative pathologies and is harmful to cells (11). High-dose vitamin E can induce nausea, headache, fatigue, double-vision, muscular weakness and kidney issues in the central nervous system (12). Many studies on vitamin E note the effects of this anti-oxidant on brain tissues and memory, as it has been seen that it improves the brain-derived neurotrophic factor by antagonising some harmful compounds (13,14). Furthermore, vitamin E in the CNS prevents brain tissue oxidative damage (15).

Vitamin E, or tocopherol, is a fat-soluble vitamin with anti-oxidant properties, fights free radicals, protects the body from the damage of pollution and cigarette smoke and protects against the onset of cancer (16). Peroxidation is a reaction involved in ageing processes and other pathologies, such as neurodegenerative ones and is harmful to cells (17). On the other hand, specific vitamins such as vitamin E are neuroprotective and relieve neurological diseases (18). While there are no treatments to halt the progression of neurodegenerative diseases, a healthy lifestyle and a diet rich in anti-oxidants can undoubtedly improve the health and function of our cells (19). On the other hand, poor nutrition or

Received: 28 March 2021
Accepted: 29 April 2021

ISSN: 2279-5855

Copyright © by BIOLIFE 2021

This publication and/or article is for individual use only and may not be further reproduced without written permission from the copyright holder. Unauthorized reproduction may result in financial and other penalties Disclosure: all authors report no conflicts of interest relevant to this article.

malnutrition can cause metabolic and nervous system disorders (20). To date, n. 8 types of vitamins E, 4 tocopherols (a, b, g, and d) and 4 tocotrienols (a, b, g, and d) have been found in nature. Vitamin E is synthesised by plants that contain alpha-tocopherol in their leaves and is extracted as an oil (21).

Alpha-tocopherol comes from the diet, is not produced by our body, and is a fat-soluble molecule. Vitamin E administration is absorbed from the small intestine and is important for the diet (22). In experimental animals, it has been seen that vitamin E taken in high doses can damage bone mineralisation, reduce hepatic storage of vitamin A, and cause coagulopathy. Several lines of evidence support that vitamin E circulates with chylomicrons and participates in the metabolism of VLDL (very low-density lipoprotein), LDL (low-density lipoprotein) and HDL (high-density lipoprotein). Mice genetically lacking high-density lipoprotein (HDL) have lower alpha-tocopherol levels and impaired brain function, demonstrating that vitamin E is closely related to HDL. In addition, alpha-tocopherol can cross the endothelial cells and arrive at the brain's glial cells, which are expressed in different brain regions and made available.

It has been seen that vitamin E is essential for the health of neurons, even if its transport and regulation in the brain remain unclear at the moment (23). Vitamin E deficiency has been observed in several neurological diseases. After being taken, it crosses the blood-brain barrier, guaranteeing a fundamental nutritional contribution to the CNS.

Vitamin E is involved in ataxia, oxidative stress, Alzheimer's disease and Parkinson's disease, where levels are low. It is unclear whether these diseases can improve after administering vitamin E (24).

Vitamin E modulates endothelial cells' function in the brain and regulates immune cell migration. Vitamin E administration is absorbed from the small intestine and is important for the diet. When given in high doses to experimental animals, a dysfunction of bone mineralisation has been observed, and reused hepatic storage of vitamin A can also give a coagulopathy. Vitamin E has excellent anti-oxidant effects with action on both inflammation and the immune system by inhibiting the effects of TNF and cellular invasiveness. Vitamin E protects against low-density lipoprotein (LDL) oxidation and promotes anti-inflammatory responses by binding to protein kinase C α by decreasing NF- κ B activation and reducing pro-inflammatory cytokine and chemokine synthesis. This effect can occur in the brain of patients with neurological diseases. The accumulation of oxidative stress raises the levels of ROS, which damage the mitochondria and counteract the cellular anti-oxidant defence mechanisms, damaging the metabolism of neurons and cerebral energy deficit. The lack of energy due to the mitochondria dysfunction leads to cell death (apoptosis), caused by an increase in the proapoptotic protein Bcl-2 Bax and Bak in the mitochondrial membrane (Fig 1).




Fig. 1. ROS are generated by mitochondria, but can also be produced by them under stress. The production of ROS causes lipid peroxidation and inositol trisphosphate stimulation, ROS activate PARPs which inhibit mitochondrial glucose with neuronal damage.

Many studies report that vitamin E, an anti-oxidant capable of eliminating ROS, is neuroprotective against cellular neurodegeneration. ROS are implicated in brain damage and stroke and can suppress the anti-oxidant defences exerted by vitamin E, causing cell death, apoptosis and necrosis. Many studies on vitamin E concern the effect of this anti-oxidant on brain tissues and also the effect on memory, as it has been seen that it improves the brain-derived neurotrophic factor by decreasing some harmful compounds (13).

CONCLUSION

In light of these observations, we can undoubtedly say that the daily intake of vitamin E at recommended doses can represent a preventive strategy against brain diseases, impaired-cell-mediated immunity and ageing and may alleviate the pathological state of the patient affected by cellular degeneration. On the other hand, low vitamin E levels with reduced alpha-tocopherol intake can lead to long-term central nervous system damage, and treatment of neurological patients deficient in vitamin E should be considered. However, further studies are required to establish the exact function of vitamin E in the brain (Table I).

Table I. Deficiencies of some vitamins that can cause disorders to the central nervous system (CNS).

VITAMIN DEFICIENCY	CNS DISORDERS
Vitamin E	encephalopathy, nerve degeneration
Riboflavin	nerve degeneration
Thiamin	peripheral neuropathy anxiety, psychosis
Niacin	depression, dementia, dizziness, irritability, tumors
Vitamin D	tetany, demineralisation, bone deformation
Vitamin A	ataxia
Vitamin B12	irritability, peripheral neuropathy, nerve degeneration

REFERENCES

1. Van Vleet, J.F.; Ferrans, V.J. Etiologic Factors and Pathologic Alterations in Selenium-Vitamin E Deficiency and Excess in Animals and Humans. *Biological Trace Element Research* 1992, *33*, 1–21, doi:10.1007/bf02783988.
2. Maltz, A. Casimer Funk, Nonconformist Nomenclature, and Networks Surrounding the Discovery of Vitamins. *The Journal of Nutrition* 2013, *143*, 1013–1020, doi:10.3945/jn.112.171827.
3. Uebanso, T.; Shimohata, T.; Mawatari, K.; Takahashi, A. Functional Roles of B-Vitamins in the Gut and Gut Microbiome. *Molecular Nutrition & Food Research* 2020, *64*, 2000426, doi:10.1002/mnfr.202000426.

4. Tyśkiewicz, K.; Dębczak, A.; Gieysztor, R.; Szymczak, T.; Rój, E. Determination of Fat- and Water-Soluble Vitamins by Supercritical Fluid Chromatography: A Review. *Journal of Separation Science* 2017, *41*, 336–350, doi:10.1002/jssc.201700598.
5. Franzke B, Schober-Halper B, Hofmann M, et al. Fat Soluble Vitamins in Institutionalized Elderly and the Effect of Exercise, Nutrition and Cognitive Training on Their Status-The Vienna Active Aging Study (VAAS): A Randomized Controlled Trial. *Nutrients*. 2019;11(6):1333. 2019. doi:10.3390/nu11061333
6. Chawla, J.; Kvarnberg, D. Hydrosoluble Vitamins. *Handbook of clinical neurology* 2014, *120*, 891–914, doi:10.1016/B978-0-7020-4087-0.00059-0.
7. Said, H.M. Water-Soluble Vitamins. *Nutrition for the Primary Care Provider* 2014, *111*, 30–37, doi:10.1159/000362294.
8. Kirilenko VN, Gregoriadis G. Fat soluble vitamins in liposomes: studies on incorporation efficiency and bile salt induced vesicle disintegration. *J Drug Target*. 1993;1(4):361-368. doi:10.3109/10611869308996095
9. Fabisiak, N.; Fabisiak, A.; Watala, C.; Fichna, J. Fat-Soluble Vitamin Deficiencies and Inflammatory Bowel Disease. *Journal of Clinical Gastroenterology* 2017, *51*, 878–889, doi:10.1097/MCG.0000000000000911.
10. McCully, K.S. Chemical Pathology of Homocysteine VIII. Effects of Tocotrienol, Geranylgeraniol, and Squalene on Thioretinaco Ozonide, Mitochondrial Permeability, and Oxidative Phosphorylation in Arteriosclerosis, Cancer, Neurodegeneration and Aging. *Annals of Clinical and Laboratory Science* 2020, *50*, 567–577.
11. Angelova, P.R.; Esteras, N.; Abramov, A.Y. Mitochondria and Lipid Peroxidation in the Mechanism of Neurodegeneration: Finding Ways for Prevention. *Medicinal Research Reviews* 2020, *41*, 770–784, doi:10.1002/med.21712.
12. Hsieh, C.-C.; Lin, B.-F. Opposite Effects of Low and High Dose Supplementation of Vitamin E on Survival of MRL/Lpr Mice. *Nutrition* 2005, *21*, 940–948, doi:10.1016/j.nut.2004.11.021.
13. Age-Related Eye Disease Study Research Group A Randomized, Placebo-Controlled, Clinical Trial of High-Dose Supplementation with Vitamins c and E, Beta Carotene, and Zinc for Age-Related Macular Degeneration and Vision Loss: AREDS Report No. 8. *Archives of ophthalmology (Chicago, Ill.: 1960)* 2001, *119*, 1417–1436, doi:10.1001/archoph.119.10.1417.
14. McCleery J, Abraham RP, Denton DA, et al. Vitamin and mineral supplementation for preventing dementia or delaying cognitive decline in people with mild cognitive impairment. *Cochrane Database Syst Rev*. 2018;11(11):CD011905. Published 2018 Nov 1. doi:10.1002/14651858.CD011905.pub2
15. Dong, S.; Huang, X.; Zhen, J.; Van Halm-Lutterodt, N.; Wang, J.; Zhou, C.; Yuan, L. Dietary Vitamin E Status Dictates Oxidative Stress Outcomes by Modulating Effects of Fish Oil Supplementation in Alzheimer Disease Model APPswe/PS1dE9 Mice. *Molecular Neurobiology* 2018, *55*, 9204–9219, doi:10.1007/s12035-018-1060-6.
16. Lee, G.; Han, S. The Role of Vitamin E in Immunity. *Nutrients* 2018, *10*, 1614, doi:10.3390/nu10111614.
17. Mazzanti, C.M.; Spanevello, R.; Ahmed, M.; Pereira, L.B.; Gonçalves, J.F.; Corrêa, M.; Schmatz, R.; Stefanello, N.; Leal, D.B.R.; Mazzanti, A.; et al. Pre-Treatment with Ebselen and Vitamin E Modulate Acetylcholinesterase Activity: Interaction with Demyelinating Agents. *International Journal of Developmental Neuroscience* 2008, *27*, 73–80, doi:10.1016/j.ijdevneu.2008.09.005.
18. Elkouzi A, Vedam-Mai V, Eisinger RS, Okun MS. Emerging therapies in Parkinson disease - repurposed drugs and new approaches. *Nat Rev Neurol*. 2019;15(4):204-223. doi:10.1038/s41582-019-0155-7
19. Finamore, A.; Palmery, M.; Bensehaila, S.; Peluso, I. Antioxidant, Immunomodulating, and Microbial-Modulating Activities of the Sustainable and Ecofriendly Spirulina. *Oxidative Medicine and Cellular Longevity* 2017, *2017*, 3247528, doi:10.1155/2017/3247528.
20. Moszak, M.; Szulińska, M.; Bogdański, P. You Are What You Eat-the Relationship between Diet, Microbiota, and Metabolic Disorders-A Review. *Nutrients* 2020, *12*, doi:10.3390/nu12041096.
21. Jensen, S.K.; Lauridsen, C. α -Tocopherol Stereoisomers. *Vitamin E* 2007, *76*, 281–308, doi:10.1016/s0083-6729(07)76010-7.
22. Hanna RM, Ghobry L, Wassef O, Rhee CM, Kalantar-Zadeh K. A Practical Approach to Nutrition, Protein-Energy Wasting, Sarcopenia, and Cachexia in Patients with Chronic Kidney Disease. *Blood Purif*. 2020;49(1-2):202-211. doi:10.1159/000504240
23. Lee, P.; Ulatowski, L.M. Vitamin E: Mechanism of Transport and Regulation in the CNS. *IUBMB Life* 2018, *71*, 424–429, doi:10.1002/iub.1993.
24. Bolotta, A.; Pini, A.; Abruzzo, P.M.; Ghezzi, A.; Modesti, A.; Gamberi, T.; Ferreri, C.; Bugamelli, F.; Fortuna, F.; Vertuani, S.;

- et al. Effects of Tocotrienol Supplementation in Friedreich's Ataxia: A Model of Oxidative Stress Pathology. *Experimental Biology and Medicine* 2020, *245*, 201–212, doi:10.1177/1535370219890873.
25. Khanna, S.; Roy, S.; Slivka, A.; Craft, T.K.S.; Chaki, S.; Rink, C.; Notestine, M.A.; DeVries, A.C.; Parinandi, N.L.; Sen, C.K. Neuroprotective Properties of the Natural Vitamin E α -Tocotrienol. *Stroke; a journal of cerebral circulation* 2005, *36*, 2258–2264, doi:10.1161/01.STR.0000181082.70763.22.