



IMMUNITY AND CANCER: IS THE VACCINATION READY FOR USE?

E. Toniato*

Department of Innovative Technologies in Medicine and Dentistry, University “G. D’Annunzio”, Chieti, Italy.

*Correspondence to:

Elena Toniato, MD,
Department of Innovative Technologies in Medicine and Dentistry,
University “G. D’Annunzio”,
Chieti, Italy.
e-mail: elena.toniato@unich.it

KEYWORDS: *immunity, cancer, vaccination, immunotherapy, tumor, inflammation*

INTRODUCTION

Cancer is a global disease and a leading cause of death worldwide, second only to cardiovascular disease in Western countries (1). Tumor cells replicate irregularly and form metastasis which tend to invade surrounding and distant tissues. In recent years, there has been a continuous evolution in the field of immunotherapy against tumors. This is also due to new diagnostic techniques and investigations such as positron emission tomography (PET), computed tomography, and magnetic resonance imaging (MRI). Immunotherapy is a therapeutic route often used by researchers that utilizes molecular investigations both for the diagnosis and treatment of tumors.

Immunotherapy makes use of the knowledge of natural killer cells (NK), CD3, CD8, and of the study of Chimeric Antigen Receptor (CAR) T cells, a type of immune cell (now the focus of many laboratories) which utilizes a patient’s T lymphocytes which are then genetically modified in a laboratory to allow them bind to cancer cells, attacking the cancer (2). For example, it has been seen that some types of oncolytic viruses destroy tumor cells without killing healthy tissue cells, inducing anti-tumor immune responses with a promising therapeutic strategy (3). The anti-cancer vaccination is making its way in small steps and is based on the use of target antigens, even if these appear very weak and of low immunogenicity. This makes it difficult to induce clinical responses and address the problem of therapeutic tumor vaccination.

Researchers are working to develop therapeutic vaccines against cancer. The use of vaccine vectors, both biological and synthetic, can improve the responses of the immune system regarding T cells, as well as B-lymphocytes and antigen-presenting cells (APC) such as macrophages and dendritic cells (4). These therapeutic strategies act on the mechanisms used by tumor cells to evade and inhibit the immune system. To date, several types of vaccines against tumors have been developed, such as antibodies that attack tumor cells, proteins, peptides, RNA, DNA, and antigens, however these are still under investigation (5). In the wake of the mRNA vaccine used against COVID-19, vaccines of this type have also been used as a new method against tumors (6). These vaccines are based on the inoculation of mRNA which encodes the production of specific tumor antigens by the host with triggering of the immune response (7). By isolating tumor mRNA, this therapy can be adapted to each type of cancer. But before these vaccines can be employed in the clinic, many challenges must be overcome, such as identifying the route of administration, mRNA efficacy, specificity, and side effects. Vaccine immunotherapy against tumors has proved to be encouraging, promising, and safe, but phase III results are still needed for the application of these vaccines in clinical oncology.

Received: 18 February 2023
Accepted: 24 March 2023

2974-6345 (2023)

Copyright © by BIOLIFE

This publication and/or article is for individual use only and may not be further reproduced without written permission from the copyright holder. Unauthorized reproduction may result in financial and other penalties. Disclosure: all authors report no conflicts of interest relevant to this article.

DISCUSSION

Lethal brain tumors have poor immune responses and are difficult to treat as they produce substances that escape the immune system (8,9). Targeting tumor antigens could reduce the development of the tumor and decrease immunosuppression. Microglia and infiltrating cells such as monocytes and lymphocytes can fuel tumor cell replication by producing pro-inflammatory and immunosuppressive cytokines (10). Targeting the tissue that provides nourishment and support of a malignant tumor, such as vascularized connective tissue and infiltrated immune cells, could be of therapeutic help. However, this stromal system, on one hand, constitutes a defense reaction of the organism to fight cancer; while on the other, it allows the tumor to develop by supplying it with the elements necessary for growth (11). Non-immune cells also contribute to the immunosuppressive environment by producing different compounds. The development of therapeutic vaccines against cancer are engaging many researchers around the world. Although cancer cells evade and suppress the immune system, it is hoped that in the future cancer vaccinations can create effective and side-effect-free anti-tumor immunity.

To date, many types of cancer vaccines have been tested which target a specific type of antigen, but so far none have been completely satisfactory (12). For example, astrocytes are cells that produce factors that feed the growth of tumor cells and favor metastasis, while neurons favor the proliferation of tumor cells by generating molecules such as neuroligin-3, a protein encoded by the NLGN3 gene that can act as a specific ligand of junction site for beta-neurexins and may be involved in central nervous system (CNS) synapses.

In recent decades, cancer therapies have had a significant improvement, also in terms of survival (13). However, despite new treatments, this effect does not apply to brain tumor therapies, which statistically, do not improve survival rates. Neoantigens are specific to the tumor and are generated by expressed gene mutations of the tumor cells (14). Therefore, personalized neoantigen vaccines can be created for therapy against certain types of tumors (15). Neoantigens can generate immune responses causing tumor rejection. In addition, neoantigenic (multi-epitope) vaccination is used for some cancers, including glioblastoma which usually has a relatively low number of mutations. This type of vaccination can stimulate neoantigen-specific CD4+ and CD8+ T lymphocytes, positively modifying the immune system of patients with glioblastoma (16). This method marks a new, encouraging step towards the immunotherapy of brain tumors.

CONCLUSIONS

In conclusion, the progress in the field of tumor immunotherapy and in vaccines against tumor antigens, is a promising new frontier in the battle against cancer. However, the results obtained thus far are not satisfactory and therefore more studies are needed to improve outcomes for this important disease which afflicts the worldwide population.

Conflict of interest

The author declares that they have no conflict of interest.

REFERENCES

1. Wang H, Naghavi M, Allen C, et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*. 2016;388(10053):1459-1544. doi:[https://doi.org/10.1016/s0140-6736\(16\)31012-1](https://doi.org/10.1016/s0140-6736(16)31012-1)
2. Mirzaei HR, Rodriguez A, Shepphird J, Brown CE, Badie B. Chimeric Antigen Receptors T Cell Therapy in Solid Tumor: Challenges and Clinical Applications. *Frontiers in Immunology*. 2017;8. doi:<https://doi.org/10.3389/fimmu.2017.01850>
3. Macedo N, Miller DM, Haq R, Kaufman HL. Clinical landscape of oncolytic virus research in 2020. *Journal for ImmunoTherapy of Cancer*. 2020;8(2):e001486. doi:<https://doi.org/10.1136/jitc-2020-001486>
4. Roth GA, Picece VCTM, Ou BS, Luo W, Pulendran B, Appel EA. Designing spatial and temporal control of vaccine responses. *Nature Reviews Materials*. 2022;7(3):174-195. doi:<https://doi.org/10.1038/s41578-021-00372-2>
5. Jahanafrooz Z, Baradaran B, Mosafer J, et al. Comparison of DNA and mRNA vaccines against cancer. *Drug Discovery Today*. 2020;25(3):552-560. doi:<https://doi.org/10.1016/j.drudis.2019.12.003>

6. Huff AL, Jaffee EM, Zaidi N. Messenger RNA vaccines for cancer immunotherapy: progress promotes promise. *Journal of Clinical Investigation*. 2022;132(6). doi:<https://doi.org/10.1172/jci156211>
7. Liu J, Fu M, Wang M, Wan D, Wei Y, Wei X. Cancer vaccines as promising immuno-therapeutics: platforms and current progress. *Journal of Hematology & Oncology*. 2022;15(1). doi:<https://doi.org/10.1186/s13045-022-01247-x>
8. Sampson JH, Gunn MD, Fecci PE, Ashley DM. Brain immunology and immunotherapy in brain tumours. *Nature Reviews Cancer*. 2019;20(1):12-25. doi:<https://doi.org/10.1038/s41568-019-0224-7>
9. Ostrom QT, Gittleman H, Liao P, et al. CBTRUS Statistical Report: Primary brain and other central nervous system tumors diagnosed in the United States in 2010–2014. *Neuro-Oncology*. 2017;19(suppl_5):v1-v88. doi:<https://doi.org/10.1093/neuonc/nox158>
10. Roesch S, Rapp C, Dettling S, Herold-Mende C. When Immune Cells Turn Bad—Tumor-Associated Microglia/Macrophages in Glioma. *International Journal of Molecular Sciences*. 2018;19(2):436. doi:<https://doi.org/10.3390/ijms19020436>
11. Seager RJ, Hajal C, Spill F, Kamm RD, Zaman MH. Dynamic interplay between tumour, stroma and immune system can drive or prevent tumour progression. *Convergent Science Physical Oncology*. 2017;3(3):034002. doi:<https://doi.org/10.1088/2057-1739/aa7e86>
12. Buonaguro L, Tagliamonte M. Selecting Target Antigens for Cancer Vaccine Development. *Vaccines*. 2020;8(4):615. doi:<https://doi.org/10.3390/vaccines8040615>
13. Miller KD, Nogueira L, Devasia T, et al. Cancer treatment and survivorship statistics, 2022. *CA: A Cancer Journal for Clinicians*. 2022;72(5). doi:<https://doi.org/10.3322/caac.21731>
14. Srivastava PK. Neoepitopes of Cancers: Looking Back, Looking Ahead. *Cancer Immunology Research*. 2015;3(9):969-977. doi:<https://doi.org/10.1158/2326-6066.cir-15-0134>
15. Liu Z, Lv J, Dang Q, et al. Engineering neoantigen vaccines to improve cancer personalized immunotherapy. *International Journal of Biological Sciences*. 2022;18(15):5607-5623. doi:<https://doi.org/10.7150/ijbs.76281>
16. Keskin DB, Anandappa AJ, Sun J, et al. Neoantigen vaccine generates intratumoral T cell responses in phase Ib glioblastoma trial. *Nature*. 2018;565(7738):234-239. doi:<https://doi.org/10.1038/s41586-018-0792-9>